

NOTES ON THE SECTION CONJECTURE OF GROTHENDIECK

FENG-WEN AN

ABSTRACT. In this short note, we will give the key point of the section conjecture of Grothendieck, that is reformulated by monodromy actions. Here, we will also give the result of the section conjecture for algebraic schemes over a number field.

Contents

Introduction

1. Preliminaries

2. Section Conjecture: Key Point

3. Section Conjecture, II: Algebraic Schemes

References

1. PRELIMINARIES

1.1. Notation. In this note, an **algebraic K -variety** is an integral scheme over a number field K of finite type.

An **arithmetic variety** is an integral scheme X satisfying the conditions:

- $\dim X \geq 1$.
- There is a surjective morphism $f : X \rightarrow \operatorname{Spec}(\mathbb{Z})$ of finite type.

For an integral scheme Z , set

- $k(Z) \triangleq$ the function field of an integral scheme Z ;
- $\pi_1^{et}(Z) \triangleq$ the étale fundamental group of Z for a geometric point of Z over a separable closure of the function field $k(Z)$.

In particular, for a field L , we set

$$\pi_1^{et}(L) \triangleq \pi_1^{et}(\operatorname{Spec}(L)).$$

2000 *Mathematics Subject Classification.* Primary 14F35; Secondary 11G35.

Key words and phrases. anabelian geometry, arithmetic scheme, étale fundamental group, section conjecture.

1.2. Outer homomorphisms. Let G, H, π_1, π_2 be four groups with homomorphisms $p : G \rightarrow \pi_1$ and $q : H \rightarrow \pi_2$, respectively.

The **outer homomorphism set** from G into H over π_1 and π_2 respectively, denoted by $Hom_{\pi_1, \pi_2}^{out}(G, H)$, is the set of the maps σ from the quotient $\frac{\pi_1}{p(G)}$ into the quotient $\frac{\pi_2}{q(H)}$ given by a group homomorphism $f : G \rightarrow H$ in such a manner:

$$\sigma : \frac{\pi_1}{p(G)} \rightarrow \frac{\pi_2}{q(H)}, x \cdot p(G) \mapsto f(x) \cdot q(H)$$

for any $x \in \pi_1$.

Remark 1.1. Let $Out(G)$ and $Out(H)$ be the outer automorphism groups. Suppose that G and H are normal subgroups of π_1 and π_2 , respectively. Then $Hom_{\pi_1, \pi_2}^{out}(G, H)$ can be regarded as a subset of $Hom(Out(G), Out(H))$.

1.3. Universal Cover and sp -Completion. For details, see [4]-[9].

Definition 1.2. Let X be an arithmetic variety (or algebraic K -variety, respectively). Then there is an integral variety $X_{\Omega_{et}}$ and a surjective morphism $p_X : X_{\Omega_{et}} \rightarrow X$ satisfying the conditions:

- $k(X_{\Omega_{et}}) = k(X)^{un}$ (or $= k(X)^{au}$, respectively);
- p_X is affine;
- $k(X_{\Omega_{et}})$ is Galois over $k(X)$;
- $X_{\Omega_{et}}$ is quasi-galois closed over X by p_X .

The integral variety $X_{\Omega_{et}}$ is called a **universal cover** over X for the étale fundamental group $\pi_1^{et}(X)$, denoted by $(X_{\Omega_{et}}, p_X)$.

Definition 1.3. For any integral variety X , there exists an integral variety X_{sp} and a surjective morphism $\lambda_X : X_{sp} \rightarrow X$ such that

- λ_X is affine;
- X_{sp} is sp -complete;
- $k(X_{sp})$ is an algebraic closure of $k(X)$;
- X_{sp} is quasi-galois closed over X by λ_X .

The integral scheme X_{sp} , is said to be an **sp -completion** of X , denoted by (X_{sp}, λ_X) .

1.4. Formally unramified extensions. (See [7, 10]). Let us recall the definition for formally unramified extension of a given field.

Definition 1.4. Let K_1 and K_2 be two arbitrary extensions over a field K such that $K_1 \subseteq K_2$.

(i) K_2 is said to be a **finite formally unramified Galois** extension of K_1 if there are two algebraic varieties X_1 and X_2 and a surjective morphism $f : X_2 \rightarrow X_1$ such that

- $k(X_1) = K_1, k(X_2) = K_2$;
- X_2 is a finite étale Galois cover of X_1 by f .

(ii) K_2 is said to be a **finite formally unramified** extension of K_1 if there is a field K_3 over K such that K_2 is contained in K_3 and K_3 is a finite formally unramified Galois extension of K_1 .

(iii) K_2 is said to be a **formally unramified** extension of K_1 if the field $K_1(\omega)$ is a finite formally unramified extension of K_1 for each element $\omega \in K_2$. In such a case, the element ω is said to be **formally unramified** over K_1 .

Let L be an arbitrary extension over a field K . Set

- $L^{al} \triangleq$ an algebraical closure of L ;
- $L^{au} \triangleq$ the maximal finite formally unramified subextensions over L contained in L^{al} ;
- $G(L) \triangleq$ the absolute Galois group $Gal(L^{al}/L)$;
- $G(L)^{au} \triangleq$ the Galois group $Gal(L^{au}/L)$ of the maximal formally unramified extension L^{au} of L .

Remark 1.5. It is seen that for the case of an algebraic extension, in general, the formally unramified extension defined in *Definition 1.4* does not coincide with that in algebraic number theory.

Remark 1.6. Note that we define another unramified extensions in [5, 6, 9] for arithmetic schemes, which is a generalization of unramified extensions in algebraic number theory and hence is different from the above one defined in *Definition 1.4*.

2. SECTION CONJECTURE: KEY POINT

The key point of the section conjecture, or what Grothendieck tells to us indeed, can be reformulated in a manner such as the following.

Theorem 2.1. *Let X, Y be two algebraic K -varieties or two arithmetic varieties. Then there exists a bijection*

$$Hom(X, Y) \cong Hom\left(\frac{Aut(X_{sp}/X)}{Aut(X_{\Omega_{et}}/X)}, \frac{Aut(Y_{sp}/Y)}{Aut(Y_{\Omega_{et}}/Y)}\right)$$

between sets.

This result is immediate from the monodromy actions of automorphism groups on the integral schemes that are considered. See [9, 10] for the proof.

Here, we have the following group isomorphisms

$$Gal(k(X_{sp})/k(X)) \cong Aut(X_{sp}/X);$$

$$\begin{aligned}
& Gal(k(Y_{sp})/k(Y)) \cong Aut(Y_{sp}/Y); \\
& \pi_1^{et}(X) \cong Gal(k(X_{\Omega_{et}})/k(X)) \cong Aut(X_{\Omega_{et}}/X); \\
& \pi_1^{et}(Y) \cong Gal(k(Y_{\Omega_{et}})/k(Y)) \cong Aut(Y_{\Omega_{et}}/Y)
\end{aligned}$$

in virtue of the properties of quasi-galois closed schemes.

3. SECTION CONJECTURE, II: ALGEBRAIC SCHEMES

For anabelian geometry of algebraic schemes over K , we have the following results. See [10] for detail.

Theorem 3.1. *Let X and Y be two algebraic K -varieties such that $k(Y)$ is contained in $k(X)$. Then there is a bijection*

$$Hom(X, Y) \cong Hom_{\pi_1^{et}(k(X)), \pi_1^{et}(k(Y))}^{out}(\pi_1^{et}(X), \pi_1^{et}(Y))$$

between sets.

Theorem 3.2. *For any algebraic K -variety X , there is a bijection*

$$\Gamma(X/K) \cong Hom_{\pi_1^{et}(K), \pi_1^{et}(k(X))}^{out}(\pi_1^{et}(K), \pi_1^{et}(X))$$

between sets.

Using Galois groups of fields, we will have the following versions of the main theorems above.

Theorem 3.3. *Let X and Y be two algebraic K -varieties such that $k(Y)$ is contained in $k(X)$. Then there is a bijection*

$$Hom(X, Y) \cong Hom_{G(k(X)), G(k(Y))}^{out}(G(k(X))^{au}, G(k(Y))^{au})$$

between sets.

Theorem 3.4. *For any algebraic K -variety X , there is a bijection*

$$\Gamma(X/K) \cong Hom_{G(K), G(k(X))}^{out}(G(K)^{au}, G(k(X))^{au})$$

between sets.

REFERENCES

- [1] An, F-W. The combinatorial graph of a scheme. eprint arXiv:0801.2609.
- [2] An, F-W. Automorphism groups of quasi-galois closed arithmetic schemes. eprint arXiv:0907.0842.
- [3] An, F-W. Automorphism groups of quasi-galois closed arithmetic schemes, (without affine structures). preprint.
- [4] An, F-W. On the existence of geometric models for function fields in several variables. eprint arXiv:0909.1993.
- [5] An, F-W. on the étale fundamental groups of arithmetic schemes, revised. eprint arXiv:0910.4646.
- [6] An, F-W. On the arithmetic fundamental groups. eprint arXiv:0910.0605.
- [7] An, F-W. On the algebraic fundamental groups. eprint arXiv:0910.4691.
- [8] An, F-W. Notes on the quasi-galois closed schemes. eprint arXiv:0911.1073.
- [9] An, F-W. On the section conjecture of Grothendieck. eprint arXiv:0911.1523.
- [10] An, F-W. On the section conjecture of Grothendieck, II. preprint.
- [11] Freitag, E; Kiehl, R. Étale Cohomology and the Weil Conjecture. Springer, Berlin, 1988.
- [12] Grothendieck, A; Dieudonné, J. Éléments de Géométrie Algébrique. vols I-IV, Pub. Math. de l'IHES, 1960-1967.
- [13] Grothendieck, A; Raynaud, M. Revêtements Étales et Groupe Fondamental (SGA1). Springer, New York, 1971.
- [14] Grothendieck, A. Letter to Faltings, in *Geometric Galois Actions*, Vol 1, edited by Schneps, L ; Lochak, P. Cambridge University Press, New York, 1997.
- [15] Hartshorne, R. Algebraic Geometry. Springer, New York, 1977.
- [16] Milne, J. Étale Cohomology. Princeton University Press, Princeton, New Jersey, 1980.
- [17] Pop, F. Glimpses of Grothendieck's anabelian geometry, in *Geometric Galois Actions*, Vol 1, edited by Schneps, L ; Lochak, P. Cambridge University Press, New York, 1997.
- [18] Suslin, A; Voevodsky, V. Singular homology of abstract algebraic varieties. Invent. Math. 123 (1996), 61-94.
- [19] Suslin, A; Voevodsky, V. Relative cycles and Chow sheaves, in *Cycles, Transfers, and Motivic Homology Theories*, Voevodsky, V; Suslin, A; Friedlander, E M. Annals of Math Studies, Vol 143. Princeton University Press, Princeton, NJ, 2000.
- [20] <http://people.math.jussieu.fr/~leila/grothendieckcircle/letters.php>

SCHOOL OF MATHEMATICS AND STATISTICS, WUHAN UNIVERSITY, WUHAN,
HUBEI 430072, PEOPLE'S REPUBLIC OF CHINA

E-mail address: fwan@amss.ac.cn